JOURNAL OF BIOMEDICAL RESEARCH & ENVIRONMENTAL SCIENCES
ISSN: 2766-2276
SUBJECT AREA(S): MUSCULOSKELETAL DISORDERS | PHYSICAL THERAPY | REHABILITATION
DOI: 10.37871

REVIEW ARTICLE

The Effects of Virtual Gaming on Lower Extremity Function in Children with Cerebral Palsy: A Narrative Review

Washington M[1], Owens RA[2] and Moody JA[1,3]*

[1]Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, Wales, United Kingdom
[2]SDR Rehabilitation UK Ltd., Cwmbran, Wales, United Kingdom
[3]School of Physical Education and Sports, Nisantasi University, Istanbul, Turkey

ABSTRACT

Background: Investigations into the use of virtual gaming, focussing on its effect on lower extremity function within therapy of children with cerebral palsy is conspicuous by its absence.

Purpose: The aim of this review is to provide an evidence-based account and associated discussion surrounding current research in the area.

Data sources: A literature search of 'PubMed', 'Google Scholar' and 'MetSearch' with relevant keywords, with article inclusion requiring publication between 2011 and 2021 was completed.

Content: Virtual reality influences multiple lower extremity functions when compared to conventional therapy or no therapy. However, information surrounding optimal prescription and participant characteristics is not evident.

Implications: Virtual gaming for use concurrently with other therapies is recommended, with the requirement for further high-quality research to advance the use of virtual gaming further and individual dose response.

***Corresponding author**

Moody JA, Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, Wales, United Kingdom

E-mail: jmoody@cardiffmet.ac.uk

DOI: 10.37871/jbres1512

Submitted: 10 July 2022

Accepted: 16 July 2022

Published: 18 July 2022

Copyright: © 2022 Washington M, et al. Distributed under Creative Commons CC-BY 4.0

OPEN ACCESS

Introduction

Cerebral Palsy (CP) is a neurological disorder occurring in children due to trauma to one or more areas of the developing brain, prior to full cerebral development [1,2]. The condition of CP is recognised worldwide, and is estimated to be present in 2-3 per 1000 live births [3]. The impaired body structure and function has significant influence on the child's quality of life impacting on their independence, inclusion and participation level, where daily activities can be significantly limited. The condition may cause both physical and psychological issues [4]. Cerebral palsy is a congenital, life-long condition where continuous therapy and strength training is necessary to maintain, and/or increase functional mobility [5].

The integration of technology within rehabilitation is rapidly developing, with Virtual Reality (VR) and Active Gaming (AG) being incorporated into both clinical and everyday life of individuals with CP [6]. Virtual reality is defined as computer hardware and software allowing interactive stimulation permitting the user to engage and participate in an environment similar to real-world experiences and objects [5]. The platforms offer a greater level of sensory stimulation, motor learning and motivation for the engagement with therapy [6-8].

Previous reviews have been conducted into motor function, upper extremity function [3,6,9-13], all confirming support towards the effectiveness of the use of

GENERAL SCIENCE
MUSCULOSKELETAL DISORDERS | PHYSICAL THERAPY | REHABILITATION
VOLUME: 3 ISSUE: 7 - JULY, 2022

Check for updates
SCAN ME

How to cite this article: Washington M, Owens RA, Moody JA. The Effects of Virtual Gaming on Lower Extremity Function in Children with Cerebral Palsy: A Narrative Review. J Biomed Res Environ Sci. 2022 July 18; 3(7): 771-778. doi: 10.37871/jbres1512, Article ID: JBRES1512, Available at: https://www.jelsciences.com/articles/jbres1512.pdf

VR as an intervention. However, currently no reviews have been conducted into multiple variables which influence Lower Extremity (LE) function making the research area somewhat conspicuous by its absence. Therefore, the current review looks to address the effectiveness of VR therapy on LE function mobility, muscle strength, gait and Range of Motion (ROM) in children with CP. Due to the scope of the review discussions surrounding research methodological approaches is limited and therefore only superficial discussions are presented.

Research Methodology

The review aims to describe studies, which have used VR as a rehabilitation tool for children with CP. Consequently the research followed narrative review guidelines from two separate studies [14,15]. A PICO framework (participants, interventions, comparators, and outcomes) was used to provide clear scope of the research required for the review [16]. PICO framework: P = Children with spastic cerebral palsy; I = Virtual reality gaming; C = A comparison therapy or no therapy; O = LE function (e.g. GMFM, strength, gait, ambulation, ROM). The process of screening and analysing studies was completed out by a single researcher.

Identification and selection of studies

This review aims to capture a representative sample of existing literature. Electronic searches were conducted on the following databases: 'PUBMED', 'Google Scholar' and 'MetSearch'. The search strategy included terms with words to the effect of 'cerebral palsy', 'virtual reality', 'video games', 'virtual gaming', 'lower extremity', 'gait', 'ambulation'. A truncation technique was used to help broaden searches; the method allows for various word endings to be applied by using an asterisk (*) in search bars (e.g. game* allows results for games and gaming). Full copies of relevant studies were obtained, and their reference list screened for further research relevant to the subject area.

Eligibility criteria

The inclusion criteria for the current review was: (a) papers written in English Language, and dated between 2011-2021, (b) the target population was between birth and 18 years old, with any classification of CP, (c) the results compared VR with conventional therapy or no therapy, (d) the study reported outcome measures surrounding the LE function and performance. The studies included often measured multiple outcome measures, only measures solely relating to the LE are included (excluding any postural or balance related measures).

Discussion

Cerebral palsy

Cerebral palsy most commonly occurs during the prenatal period with 80% of cases occurring during this time, mainly due to unknown causes; however, complications such as asphyxia have been confirmed to cause CP in some cases [1]. Cerebral Palsy can also occur during postnatal and neonatal periods [17], with common risk factors including a child weighing <2.5kg at birth, premature birth, brain haemorrhage, child abuse and bacterial meningitis [1]. The damage can lead to musculoskeletal and neuromuscular impairments, ultimately effecting daily living of children and adults with CP [2].

Individuals with CP are most commonly affected by motor impairment, which is commonly displayed through muscle weakness, muscle hypertonicity, tremors, scissor-gait and increased tendon reflexes [18]. However, the disease can often be accompanied by less apparent symptoms such as disturbances to sensation, perception, cognitive and behavioural performances [2,9].

The Gross Motor Function Classification System (GMFCS) is one of many standardised assessments used to assess the severity of CP. It consists of five levels representing differences in function relating to daily life [19]. The assessment is widely accepted throughout research with a strong inter-rather reliability (0.84) for children younger than six and 0.89 for children 6 years of age and older [20]. Other scales used in research include, Child Health Questionnaire, Paediatric Evaluation of Disability Inventory, Functional Independence Measure, and Ashworth scale [21-24]. With such grading in place it allows CP patients quality of life and development milestones to be monitored effectively [2].

A range of techniques has been used to minimise and lessen the severity of CP [25-27]; however, the interventions cannot eliminate CP completely and therefore, treatments must be on-going and ever-present throughout an individual's life. With a lack of access to available therapy in the UK, especially, as patients get older, multiple time consuming appointments and expensive equipment, alternative approaches are being used to make treatment more flexible, enjoyable and affordable [28,29].

Virtual reality

An increase in technology has permitted gaming platforms and devices to be integrated into CP rehabilitation. Virtual reality therapy and active gaming have increasingly become a popular method for home, school and in-clinic rehabilitation interventions [30]. The systems can be used to achieve rehabilitation goals by immersing individuals into 3D games, whilst adapting game type, complexity, and intensity to suit the individual therapy requirements of the individual, as well as children being able to carry-out real-life activities without the risk of real-life hazards [8,31,32].

There are two versions of VR systems for rehabilitation, with the first being commercial video game platforms and devices, this includes devices such as the Nintendo Wii, Sony

PlayStation and Xbox Consoles [30]. The second method is custom VR systems which are specifically developed for rehabilitation purposes, systems such as, Move to improve it (MiTII; Mitii Development A/S) [33], E-Link Evaluation and Exercise System and Eloton SimCycle Virtual Cycling System (Eloton, Inc., NV, USA). Both commercial and custom systems offer a more enjoyable experience for users and subsequently, increase motivation and buy-in for rehabilitation purposes [34-36].

Commercial and custom systems have been reported to have positive effects in numerous LE research studies with improvements being shown in mobility, daily living ability, gait and other limb components [29,37-42]. However, no significant findings are still evident throughout the research with a variety of methodologies and equipment proving unable to elicit responses when compared with other therapies or no therapy [43]. A previous review in the area concluded that interventions were able to elicit improvements to gross motor function and skill transfer to daily living tasks and for the brain to create new connections between neurons and motor learning requirements to be exploited to make relatively permanent motor changes [44-46]. Approaches such as task repetition, task difficulty and motivation all contribute to motor-learning development [47,48]. Papers within the review suggest that motor learning can be potentiated with VR and therefore, skills can transfer to motor abilities and real-life integration [49-51]. However, the review was comprised of papers which lacked a high- quality methodology and subsequently reduces the credibility of findings [52].

A more recent meta-analysis of randomised control trials with a fair to good Pedro quality determined that VR was a viable intervention for improving ambulation. Only 8 of the 19 papers included in the review measured ambulation variables, the papers showed a medium to large effect size of 0.755. However, with the detection of publication bias a Duval-Tweedie trim and fill technique was applied which reduced the effect size to 0.378, suggesting that the findings of the results are sensitive to change [53]. In addition, the authors did not provide information on which ambulation variables of gait velocity or gait endurance were enhanced, therefore speculation remains surrounding which attributes are enhanced. Further conclusions related to engineer-built systems showing a greater effect than commercial systems, it could be suggested that specifically built systems more optimally target the child's needs with adjustments of game difficulty and training goals being available [54]. In addition, commercial platforms aren't created for rehabilitation purposes and therefore have limited therapeutic value. However, these findings are not exclusive to LE variables.

Lower extremity functions

Children with CP are faced with a variety of symptoms which can alter their overall LE function and ultimately their quality of life. Previous investigations of VR effects on LE functions have focused on a child's ability to walk, stand and additional functional movements [12,55,56]. These movements although considered to be a child's later life goals, cover a vast amount of the body of research due to the importance of such aspects and the contribution it has to for life-long rehabilitation [55,56]. However, additional components are evident throughout research and will also be discussed within the review. The review will aim to discuss research surrounding ambulation and gait, gross motor function measures, functional mobility, muscle strength and Range of Motion (ROM) of the LE. Although ambulation is considered through both functional mobility and gross motor function, it will be considered separately to ensure thorough discussions surrounding the necessary gait variables. These include, gait velocity, gait endurance, gait cadence, stride length and step length. It should also be noted that although gait and gross motor function measures are directly related to functional mobility, they are discussed separately in the review.

Virtual reality and functional mobility

Functional mobility is required to perform daily activities, to enhance participation and functional independence [43,57]. A primary goal of many CP rehabilitation programmes is to reduce limitations related to lower extremity function which influences the ability to walk, ascend and descend stairs [58]. The most recognised and commonly used method throughout CP research is the Gross Motor Function Measure [59], which will be discussed further in the review. Other LE functional tests have been used by following at 12-week intervention using the WiiFit system (Nintendo, Kyoto, Japan). Children carried out the 10-stair climbing test, where they were tested on their ability to climb 10 steps as fast as possible [60], significant improvements were observed from the VR therapy group compared to the control group, although only a small effect size was evident (p = 0.001; ES= 0.19). Moreover, the Timed Up and Go (TUG) test was also measured and findings showed improvements for both groups, with the VR therapy group yielding a significantly faster performance (p = 0.001; effect size [ES] = 0.48). It is suggested that tailored games to test balance and weight-transfer of the LE allowed children to become more stable when completing functional movements [42].

Conversely, further investigations into the TUG measure have elicited no significant differences to conventional therapies. The studies also found other non-significant differences between groups in other measures [38,61], including a timed up and down stairs measure. However, there were observed trends towards experimental groups, suggesting that the programmes provided may not have been a sufficient intensity or length to elicit responses [61].

Qualitative measures have also been used to assess the influence of VR therapy on daily living activities, the functional independence measure for children test has an

strong reliability (ICC = 0.91-0.98) and indicated significantly greater responses following a Wii Fit VBT intervention [42]. Conversely, following the use of the same gaming platform the 28-mobility measure and quantitative analysis of participants physical activity over a four-day period was unable to confirm significant differences to conventional training. Although, within the study performance capacity was shown to have greater improvements this was not replicated for activity performance or mobility limitations. Numerous barriers are associated with a child's participation in activity such as school, peers and therapy [62], therefore it is suggested that ecological influence would be evident throughout physical performance measures.

Virtual reality and gross motor function measures

Gross motor function measures are used to assess motor skills within children with cerebral palsy. The GMFM, along with other modified versions (GMFM-66, which consists of 66 items and GMFM-88, consisting of 88 items) covers the five components of lying and rolling, sitting, crawling and kneeling, standing and walking, running and jumping [63]. A meta-analysis found a medium effect size (hedges g = 0.44) of GMFM scores following VR therapy of six studies; it also revealed that a shorter programme session of 20-30 minutes length led to a greater influence on adaptation when compared to a longer session (40-45 minutes) with an effect size of 0.56 compared to 0.14 respectively [64]. However, the recommended programme prescription is limited to length of session and is limited to six study designs; therefore, speculation remains surrounding optimal prescription.

Additional VR therapy is reported to have no significant difference to conventional therapies following a 12-week intervention [6,65]. The studies only provided a whole score for the GMFM, so details regarding the influence on individual elements of standing and walking, and running and jumping which were considered in the meta-analysis, are not able available for comparison. It is important to note the research were able to elicit other lower extremity responses more specific to the VR intervention [6].

Further investigations confirmed both groups were able to elicit significant improvements for standing and walking, and running and jumping ($p < 0.05$) [38]. However, following group comparison only the latter showed a significant difference following VR therapy. The study also used a GMFM which used the same grading items but rated the quality of each movement rather than the ability to complete the movement [63]. The results showed both groups had an increase in standing variables but only the VR intervention group showed a significant increase in walking variables [38]. However, no other investigations have used the performance measure and reliability investigations suggest that highly skilled therapists are required to complete this form of analysis sufficiently [66]. Although the measure offers interesting insights surrounding the motor performance of CP children, further research is required.

Virtual reality and gait

Several methods throughout CP research have been used to assist reduce muscle over-activity and imbalance, ranging from surgery to walking aids [67]. The functional movement requires musculoskeletal integrity and neurological control between multiple joints and agonist and antagonist muscle groups simultaneously, therefore the challenges faced by children with CP can often lead to the inability to walk or gait disturbance [68,69]. A recently published meta-analysis used 16 studies to evaluate gait ability and the recommended training programme of VR therapy. All the gait sourced articles for the current review were used within the meta-analysis, the review rated the quality of methodologies using a Pedro scale [70], on average studies reported 'good' quality (5.7 mean 1.4 SD) [64]. Three gait parameters which were analysed throughout the review are confirmed to improve performance following VR intervention, this included gait velocity (hedges g = 0.76), cadence (g = 0.80), stride length (g = 0.76). Stride width was reported to have a small negative effect (g = -0.23).

These results are supported by an aforementioned review, with a 0.75 effect size being provided in ambulation function [4]. It is suggested that the results presented across the two meta-analyses could be due to different underlying mechanisms associated with VR therapy. The first is related to the therapy offering a multi-sensory (visual, audial and proprioceptive) feedback of movements, therefore allowing an enhanced retention of movement amplitudes and limb position and ultimately improving gross motor function [71-73]. Secondly, VR therapy allows participants to focus on control and movement timings of real-life activities, thus, with practice an enhanced efficiency of movement pattern and improved kinematic scores can be seen [41,74,75].

Virtual reality and lower extremity strength

Children with CP have been shown to have smaller muscles and reduced muscular strength than their healthier peers, which impedes their gross motor function and gait [33,76-80]. Mitchell investigated strength measures through maximal repetition tasks of multiple functional movements (sit-to- stand, lateral step up and half kneel) over a 30-second period [81,82].

Following a 20-week intervention the findings showed a 19.3 mean difference in the functional strength scores ($p < 0.001$). Measures into the sit-to-stand were also observed following a 12-week Wii-Fit intervention, both groups were able to show significant increases in repetitions ($p < 0.05$), but greater increases were observed for the VR group ($p < 0.001$, effect size = 0.70) [42]. The studies used different VR platforms and completely different methodological approaches but were both able to elicit function strength improvements of the LE. However, both studies noted that some parts of the protocols were unblended and therefore, could suggest potential for bias.

Further investigations compared VR treadmill training to conventional treadmill training for responses in knee extensor and flexor strength. The study reported that the VR training had been significantly more effective in improving muscular strength than treadmill training alone, with right knee flexion strength reporting no significant differences between experimental groups. It has been suggested that the VR system offers a greater reflection of daily life and therefore, the child becomes more task-orientated whilst participating [37]. Furthermore, the child will need to problem solve through repetitive movements, which facilitates an increase in motor learning, in addition, a structured level and rewards system throughout leads to a greater motivation to repeat tasks [8]. The study did have a limited sample size and therefore, can limit data saturation and accurate conclusions from the subject population.

Virtual reality and lower extremity range of movement

Both muscle shortening and decreased ROM is common within children with CP and is commonplace in patients with and without spasticity [8,83]. Contractures in the LE commonly seen through flexion at the hips and knees and plantarflexion of the ankles, subsequently affects alignments and causes imbalance around the limbs, ultimately effecting the functional ability to walk, stand and sit [84-86]. Research is conspicuous by its absence into the influence of VR therapy on LE ROM, with only three investigations assessing this function.

We measured changes to the ankle dorsiflexion angle at initial of foot contact during a gait analysis. Three participants carried out repeated active plantarflexion and dorsiflexion movements on each ankle for 20 minutes; they were not compared against a control group [87]. The results showed only one participant had significantly improved their ankle dorsiflexion, with the additional participants having little change. The study presents many limitations surrounding lack of control group and unblinding of the therapist, leaving speculation surrounding whether findings were due to the experimental procedure, influenced by external variables, or bias [88].

A study by Fowler used a Selective Control Assessment of the Lower Extremity (SCALE; ICC 0.88-0.91) tool to assess the multiple motor functions, including ankle dorsiflexion, knee extension and hip abduction [89]. The VR group and control group completed 40 minutes of conventional therapy, following this, the experimental group completed an additional 40 minutes of VR therapy. The results indicated that both groups significantly improved following the 6-week intervention, with the VR group demonstrating an increased between group difference ($p < 0.05$) across all variables except the right hip abductor. However, the measurement tool was shown to have a limiting influence on statistical changes due to a sealing effect. Therefore, it could be suggested that improvements observed for the VR group are due to the additional therapy completed [90].

Similarly, Curtis found significant improvements to passive and active ankle dorsiflexion with an extended knee ($p = 0.02$ and $p = 0.001$, respectively) following a 10-week training intervention [91]. Although, no significant changes were found for a flexed knee. Differences within the two knee positions is suggested due to the intervention being completed with an extended knee and thus, greater adaptations are caused to the gastrocnemius muscle and a greater response in ankle dorsiflexion with an extended knee. The ankle ROM was collected through the use of a modified goniometer, which shows high reliability (0.9) when used for children with CP [92]. The test was completed three times and the therapist was blinded to the individual results to eliminate measurement bias.

Further limitations were presented between the first two studies, a reduction in follow-up of participants and all studies lacked a sufficient sample size. It should also be understood that two of the studies used robotic and specialised equipment, which is not available to many clinics or patients and therefore limits the generalisation to more affordable and commercial platforms. It is clear through this section of the review that ROM improvements are observed, however, advanced methodologies with confirmed reliability are required to draw conclusions on whether improvements to ROM are solely the result of VR therapy.

Practical application

Virtual reality therapy research has shown it can progress LE functions; however, speculation remains surrounding the research and some of the experimental methodologies used. It is clear throughout the current review that VR therapy has no detrimental influence of function when used alongside other therapies [6,12,93-95]. Therefore, the use of commercial and specific platforms to improve lower extremity function is advised if available to the child. However, research is still requiring providing direction and guidance on optimal dose prescription; furthermore, clinicians should consider a patient's individual characteristics during the use of VR therapy [96].

Limitations

The narrative review structure used can be associated with author bias, with interpretation and conclusions from data extracted subjectively [97]. The review may also be influenced by selection bias, with a single reviewer selecting all studies discussed throughout the composition [98,99]. Further, the scope of the study limited the discussion surrounding the LE function topics, with important areas such as programme design and participant characteristics consider as beyond the scope of this review. Therefore, the findings of the review are limited and a more systematic review considered as an alternative for future reviews.

Conclusion

In conclusion, it is evident through the different functions that VR therapy can influence and facilitate LE performance in children with CP. The current review findings would advise VR therapy to be used in conjunction with other conventional therapies, until a more research is conducted into the sole use of VR therapy. Future research is required to review the influence of participant characteristics on results and the complex methodological approaches, which are utilised to elicit improvements in LE functions, with hope that conclusions can be drawn on optimal dose prescription for children with CP.

References

1. Krigger KW. Cerebral palsy: An overview. Am Fam Physician. 2006 Jan 1;73(1):91-100. PMID: 16417071.
2. Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, Dan B, Jacobsson B. A report: The definition and classification of cerebral palsy april 2006. Dev Med Child Neurol Suppl. 2007 Feb;109:8-14. Erratum in: Dev Med Child Neurol. 2007 Jun;49(6):480. PMID: 17370477.
3. Johansen T, Strøm V, Simic J, Rike PO. Effectiveness of training with motion-controlled commercial video games on hand and arm function in young people with cerebral palsy: A systematic review and meta-analysis. J Rehabil Med. 2019 Dec 3;52(1):jrm00012. doi: 10.2340/16501977-2633. Epub ahead of print. PMID: 31794044.
4. Chen Y, Fanchiang HD, Howard A. Effectiveness of virtual reality in children with cerebral palsy: A systematic review and meta-analysis of randomized controlled trials. Phys Ther. 2018 Jan 1;98(1):63-77. doi: 10.1093/ptj/pzx107. PMID: 29088476; PMCID: PMC6692882.
5. Weiss PL, Tirosh E, Fehlings D. Role of virtual reality for cerebral palsy management. J Child Neurol. 2014 Aug;29(8):1119-1124. doi: 10.1177/0883073814533007. Epub 2014 May 5. PMID: 24799367.
6. Chen CL, Chen CY, Liaw MY, Chung CY, Wang CJ, Hong WH. Efficacy of home-based virtual cycling training on bone mineral density in ambulatory children with cerebral palsy. Osteoporos Int. 2013 Apr;24(4):1399-1406. doi: 10.1007/s00198-012-2137-0. Epub 2012 Sep 28. PMID: 23052930.
7. Weiss PL, Rand D, Katz N, Kizony R. Video capture virtual reality as a flexible and effective rehabilitation tool. J Neuroeng Rehabil. 2004 Dec 20;1(1):12. doi: 10.1186/1743-0003-1-12. PMID: 15679949; PMCID: PMC546410.
8. Levac D, Rivard L, Missiuna C. Defining the active ingredients of interactive computer play interventions for children with neuromotor impairments: A scoping review. Res Dev Disabil. 2012 Jan-Feb;33(1):214-223. doi: 10.1016/j.ridd.2011.09.007. Epub 2011 Oct 11. PMID: 22093667.
9. Ravi DK, Kumar N, Singhi P. Effectiveness of virtual reality rehabilitation for children and adolescents with cerebral palsy: An updated evidence-based systematic review. Physiotherapy. 2017 Sep;103(3):245-258. doi: 10.1016/j.physio.2016.08.004. Epub 2016 Sep 27. PMID: 28109566.
10. Rathinam C, Mohan V, Peirson J, Skinner J, Nethaji KS, Kuhn I. Effectiveness of virtual reality in the treatment of hand function in children with cerebral palsy: A systematic review. J Hand Ther. 2019 Oct-Dec;32(4):426-434.e1. doi: 10.1016/j.jht.2018.01.006. Epub 2018 Jul 14. PMID: 30017414.
11. Wu J, Loprinzi PD, Ren Z. The rehabilitative effects of virtual reality games on balance performance among children with cerebral palsy: A meta-analysis of randomized controlled trials. Int J Environ Res Public Health. 2019 Oct 28;16(21):4161. doi: 10.3390/ijerph16214161. PMID: 31661938; PMCID: PMC6861947.
12. Ghai S, Ghai I. Virtual reality enhances gait in cerebral palsy: A training dose-response meta-analysis. Front Neurol. 2019 Mar 26;10:236. doi: 10.3389/fneur.2019.00236. PMID: 30984095; PMCID: PMC6448032.
13. Cano Porras D, Siemonsma P, Inzelberg R, Zeilig G, Plotnik M. Advantages of virtual reality in the rehabilitation of balance and gait: Systematic review. Neurology. 2018 May 29;90(22):1017-1025. doi: 10.1212/WNL.0000000000005603. Epub 2018 May 2. PMID: 29720544.
14. Green BN, Johnson CD, Adams A. Writing narrative literature reviews for peer-reviewed journals: Secrets of the trade. J Chiropr Med. 2006 Autumn;5(3):101-117. doi: 10.1016/S0899-3467(07)60142-6. PMID: 19674681; PMCID: PMC2647067.
15. Ferrari R. Writing narrative style literature reviews. Medical Writing. 2015;24(4):230-235.
16. Smith V, Devane D, Begley CM, Clarke M. Methodology in conducting a systematic review of systematic reviews of healthcare interventions. BMC Med Res Methodol. 2011 Feb 3;11(1):15. doi: 10.1186/1471-2288-11-15. PMID: 21291558; PMCID: PMC3039637.
17. Krägeloh-Mann I, Cans C. Cerebral palsy update. Brain Dev. 2009 Aug;31(7):537-544. doi: 10.1016/j.braindev.2009.03.009. Epub 2009 Apr 21. PMID: 19386453.
18. Porter RS, Kaplan JL. The Merck manual of diagnosis and therapy. Merck Sharp & Dohme Corp. 2011.
19. Palisano RJ, Avery L, Gorter JW, Galuppi B, McCoy SW. Stability of the gross motor function classification system, manual ability classification system, and communication function classification system. Dev Med Child Neurol. 2018 Oct;60(10):1026-1032. doi: 10.1111/dmcn.13903. Epub 2018 May 4. PMID: 29726578.
20. Rutz E, Tirosh O, Thomason P, Barg A, Graham HK. Stability of the gross motor function classification system after single-event multilevel surgery in children with cerebral palsy. Dev Med Child Neurol. 2012 Dec;54(12):1109-1113. doi: 10.1111/dmcn.12011. Epub 2012 Oct 16. PMID: 23067343.
21. Raat H, Botterweck AM, Landgraf JM, Hoogeveen WC, Essink-Bot ML. Reliability and validity of the short form of the child health questionnaire for parents (CHQ-PF28) in large random school based and general population samples. J Epidemiol Community Health. 2005 Jan;59(1):75-82. doi: 10.1136/jech.2003.012914. PMID: 15598731; PMCID: PMC1763365.
22. Berg N. Non-response bias. Epidemiology. 2005.
23. Linacre JM, Heinemann AW, Wright BD, Granger CV, Hamilton BB. The structure and stability of the Functional Independence Measure. Arch Phys Med Rehabil. 1994 Feb;75(2):127-132. PMID: 8311667.
24. Ashworth B. Preliminary trial of carisoprodol in multiple sclerosis. Practitioner. 1964 Apr;192:540-542. PMID: 14143329.
25. Boyd RN, Graham HK. Objective measurement of clinical findings in the use of botulinum toxin type a for the management of children with cerebral palsy. European Journal of Neurology. 1999;6(S4):23-35.
26. Novak I, Morgan C, Fahey M, Finch-Edmondson M, Galea C, Hines A, Langdon K, Namara MM, Paton MC, Popat H, Shore B, Khamis A, Stanton E, Finemore OP, Tricks A, Te Velde A, Dark L, Morton N, Badawi N. State of the evidence traffic lights 2019: Systematic review of interventions for preventing and treating children with cerebral palsy. Curr Neurol Neurosci Rep. 2020 Feb 21;20(2):3. doi: 10.1007/s11910-020-1022-z. PMID: 32086598; PMCID: PMC7035308.
27. Ritzmann R, Stark C, Krause A. Vibration therapy in patients with cerebral palsy: a systematic review. Neuropsychiatr Dis Treat. 2018 Jun 18;14:1607-1625. doi: 10.2147/NDT.S152543. PMID: 29950843; PMCID: PMC6018484.
28. Coombe S, Moore F, Bower E. A national survey of the amount of physiotherapy intervention given to children with cerebral palsy in the UK in the NHS. Association of Paediatric Chartered Physiotherapists. 2012;3(1):5-18.
29. Farr WJ, Green D, Bremner S, Male I, Gage H, Bailey S, Speller S, Colville V, Jackson M, Memon A, Morris C. Feasibility of a randomised controlled trial to evaluate home-based virtual reality therapy in children with cerebral palsy. Disabil Rehabil. 2021 Jan;43(1):85-97. doi: 10.1080/09638288.2019.1618400. Epub 2019 May 25. PMID: 31131641.
30. Demers M, Fung K, Subramanian SK, Lemay M, Robert MT. Integration of motor learning principles into virtual reality interventions for individuals with cerebral palsy: Systematic review. JMIR Serious Games. 2021 Apr 7;9(2):e23822. doi: 10.2196/23822. PMID: 33825690; PMCID: PMC8060861.
31. Burke JW, McNeill MDJ, Charles DK, Morrow PJ, Crosbie JH, Donough SM. Optimising engagement for stroke rehabilitation using serious games. The Visual Computer. 2009;25(12):1085-1099.
32. Wilson PN, Foreman N, Stanton D. Virtual reality, disability and rehabilitation. Disabil Rehabil. 1997 Jun;19(6):213-220. doi: 10.3109/09638289709166530. PMID: 9195138.
33. Mitchell LE, Ziviani J, Boyd RN. A randomized controlled trial of web-based training to increase activity in children with cerebral palsy. Dev Med Child Neurol. 2016 Jul;58(7):767-773. doi: 10.1111/dmcn.13065. Epub 2016 Feb 15. PMID: 26877078
34. Rostami HR, Arastoo AA, Nejad SJ, Mahany MK, Malamiri RA, Goharpey S. Effects of modified constraint-induced movement therapy in virtual environment on upper-limb

function in children with spastic hemiparetic cerebral palsy: a randomised controlled trial. NeuroRehabilitation. 2012;31(4):357-365. doi: 10.3233/NRE-2012-00804. PMID: 23232158.

35. Chen CL, Hong WH, Cheng HY, Liaw MY, Chung CY, Chen CY. Muscle strength enhancement following home-based virtual cycling training in ambulatory children with cerebral palsy. Res Dev Disabil. 2012 Jul-Aug;33(4):1087-1094. doi: 10.1016/j.ridd.2012.01.017. Epub 2012 Feb 28. PMID: 22502833.

36. Bryanton C, Bossé J, Brien M, McLean J, McCormick A, Sveistrup H. Feasibility, motivation, and selective motor control: virtual reality compared to conventional home exercise in children with cerebral palsy. Cyberpsychol Behav. 2006 Apr;9(2):123-128. doi: 10.1089/cpb.2006.9.123. PMID: 16640463.

37. Cho C, Hwang W, Hwang S, Chung Y. Treadmill training with virtual reality improves gait, balance, and muscle strength in children with cerebral palsy. Tohoku J Exp Med. 2016 Mar;238(3):213-218. doi: 10.1620/tjem.238.213. PMID: 26947315.

38. Rgen M, Akbayrak T, Gunel M, Cankaya O, Guchan Z, Turkyilmaz E. Investigation of the effects of the Nintendo® Wii-Fit training on balance and advanced motor performance in children with spastic hemiplegic cerebral palsy: A Randomized Controlled Trial. Int J Ther Rehabil Res. 2016;5:146.

39. James S, Ziviani J, Ware RS, Boyd RN. Randomized controlled trial of web-based multimodal therapy for unilateral cerebral palsy to improve occupational performance. Dev Med Child Neurol. 2015 Jun;57(6):530-538. doi: 10.1111/dmcn.12705. Epub 2015 Feb 27. PMID: 25955443.

40. Booth ATC, Buizer AI, Meyns P, Oude Lansink ILB, Steenbrink F, van der Krogt MM. The efficacy of functional gait training in children and young adults with cerebral palsy: A systematic review and meta-analysis. Dev Med Child Neurol. 2018 Sep;60(9):866-883. doi: 10.1111/dmcn.13708. Epub 2018 Mar 7. PMID: 29512110.

41. Gagliardi C, Turconi AC, Biffi E, Maghini C, Marelli A, Cesareo A, Diella E, Panzeri D. Immersive virtual reality to improve walking abilities in cerebral palsy: A pilot study. Ann Biomed Eng. 2018 Sep;46(9):1376-1384. doi: 10.1007/s10439-018-2039-1. Epub 2018 Apr 27. PMID: 29704186.

42. Tarakci D, Ersoz Huseyinsinoglu B, Tarakci E, Razak Ozdincler A. Effects of nintendo Wii-Fit® video games on balance in children with mild cerebral palsy. Pediatr Int. 2016 Oct;58(10):1042-1050. doi: 10.1111/ped.12942. Epub 2016 Aug 23. PMID: 26858013.

43. Sajan JE, John JA, Grace P, Sabu SS, Tharion G. Wii-based interactive video games as a supplement to conventional therapy for rehabilitation of children with cerebral palsy: A pilot, randomized controlled trial. Dev Neurorehabil. 2017 Aug;20(6):361-367. doi: 10.1080/17518423.2016.1252970. Epub 2016 Nov 15. PMID: 27846366.

44. Massetti T, Silva TDD, Ribeiro DC, Malheiros SRP, Favero FM, Monteiro. Motor learning through virtual reality in cerebral palsy–a literature review. MedicalExpress. 2014;1:302-306.

45. Krakauer JW. The applicability of motor learning to neurorehabilitation. Oxford textbook of neurorehabilitation. 2015;1:55-64.

46. Winstein CJ, Kay DB. Translating the science into practice: shaping rehabilitation practice to enhance recovery after brain damage. Prog Brain Res. 2015;218:331-360. doi: 10.1016/bs.pbr.2015.01.004. Epub 2015 Mar 19. PMID: 25890145.

47. Kleim JA, Jones TA. Principles of experience-dependent neural plasticity: Implications for rehabilitation after brain damage. J Speech Lang Hear Res. 2008 Feb;51(1):S225-239. doi: 10.1044/1092-4388(2008/018). PMID: 18230848.

48. Muratori LM, Lamberg EM, Quinn L, Duff SV. Applying principles of motor learning and control to upper extremity rehabilitation. J Hand Ther. 2013 Apr-Jun;26(2):94-102; quiz 103. doi: 10.1016/j.jht.2012.12.007. PMID: 23598082; PMCID: PMC3773509.

49. Golomb MR, Donald BC, Warden SJ, Yonkman J, Saykin AJ, Shirley B, Huber M, Rabin B, Abdelbaky M, Nwosu ME, Masih M, Burdea GC. In-home virtual reality videogame telerehabilitation in adolescents with hemiplegic cerebral palsy. Arch Phys Med Rehabil. 2010 Jan;91(1):1-8.e1. doi: 10.1016/j.apmr.2009.08.153. PMID: 20103390.

50. Grecco LA, Duarte Nde A, de Mendonça ME, Pasini H, Lima VL, Franco RC, de Oliveira LV, Carvalho T, Corrêa JC, Collange NZ, Sampaio LM, Galli M, Fregni F, Oliveira CS. Effect of transcranial direct current stimulation combined with gait and mobility training on functionality in children with cerebral palsy: Study protocol for a double-blind randomized controlled clinical trial. BMC Pediatr. 2013 Oct 11;13:168. doi: 10.1186/1471-2431-13-168. PMID: 24112817; PMCID: PMC3852945.

51. Straker LM, Campbell AC, Jensen LM, Metcalf DR, Smith AJ, Abbott RA, Pollock CM, Piek JP. Rationale, design and methods for a randomised and controlled trial of the impact of virtual reality games on motor competence, physical activity, and mental health in children with developmental coordination disorder. BMC Public Health. 2011 Aug 18;11:654. doi: 10.1186/1471-2458-11-654. PMID: 21851587; PMCID: PMC3166932.

52. Stone J, Gurunathan U, Glass K, Munn Z, Tugwell P, Doi SAR. Stratification by quality induced selection bias in a meta-analysis of clinical trials. J Clin Epidemiol. 2019 Mar;107:51-59. doi: 10.1016/j.jclinepi.2018.11.015. Epub 2018 Nov 17. PMID: 30458262.

53. Duval S, Tweedie R. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics. 2000 Jun;56(2):455-463. doi: 10.1111/j.0006-341x.2000.00455.x. PMID: 10877304.

54. Guadagnoli MA, Lee TD. Challenge point: A framework for conceptualizing the effects of various practice conditions in motor learning. J Mot Behav. 2004 Jun;36(2):212-224. doi: 10.3200/JMBR.36.2.212-224. PMID: 15130871.

55. Warnier N, Lambregts S, Port IV. Effect of virtual reality therapy on balance and walking in children with cerebral palsy: A systematic review. Dev Neurorehabil. 2020 Nov;23(8):502-518. doi: 10.1080/17518423.2019.1683907. Epub 2019 Nov 1. PMID: 31674852.

56. Damiano DL. Activity, activity, activity: Rethinking our physical therapy approach to cerebral palsy. Phys Ther. 2006 Nov;86(11):1534-1540. doi: 10.2522/ptj.20050397. PMID: 17094192.

57. Mollenkop H, Marcellini F, Ruoppila I, Flaschenträger P, Gagliardi C, Spazzafumo L. Outdoor mobility and social relationships of elderly people. Archives of gerontology and geriatrics. 1997;24(3):295-310.

58. Van Ravesteyn NT, Dallmeijer AJ, Scholtes VA, Roorda LD, Becher JG. Measuring mobility limitations in children with cerebral palsy: interrater and intrarater reliability of a mobility questionnaire (MobQues). Dev Med Child Neurol. 2010 Feb;52(2):194-199. doi: 10.1111/j.1469-8749.2009.03341.x. Epub 2009 Sep 11. PMID: 19747207.

59. Russell DJ, Rosenbaum PL, Cadman DT, Gowland C, Hardy S, Jarvis S. The gross motor function measure: a means to evaluate the effects of physical therapy. Dev Med Child Neurol. 1989 Jun;31(3):341-352. doi: 10.1111/j.1469-8749.1989.tb04003.x. PMID: 2753238.

60. Studenski S, Perera S, Wallace D, Chandler JM, Duncan PW, Rooney E, Fox M, Guralnik JM. Physical performance measures in the clinical setting. J Am Geriatr Soc. 2003 Mar;51(3):314-322. doi: 10.1046/j.1532-5415.2003.51104.x. PMID: 12588574.

61. Salem Y, Gropack SJ, Coffin D, Godwin EM. Effectiveness of a low-cost virtual reality system for children with developmental delay: A preliminary randomised single-blind controlled trial. Physiotherapy. 2012 Sep;98(3):189-195. doi: 10.1016/j.physio.2012.06.003. Epub 2012 Jul 31. PMID: 22898574.

62. Verschuren O, Wiart L, Hermans D, Ketelaar M. Identification of facilitators and barriers to physical activity in children and adolescents with cerebral palsy. J Pediatr. 2012 Sep;161(3):488-494. doi: 10.1016/j.jpeds.2012.02.042. Epub 2012 Apr 10. PMID: 22494875.

63. Boyce WF, Gowland C, Rosenbaum PL, Lane M, Plews N, Goldsmith CH, Russell DJ, Wright V, Potter S, Harding D. The gross motor performance measure: Validity and responsiveness of a measure of quality of movement. Phys Ther. 1995 Jul;75(7):603-613. doi: 10.1093/ptj/75.7.603. PMID: 7604079.

64. Ghai S, Ghai I, Lamontagne A. Virtual reality training enhances gait post stroke: A systematic review and meta-analysis. Ann N Y Acad Sci. 2020 Oct;1478(1):18-42. doi: 10.1111/nyas.14420. Epub 2020 Jul 13. Erratum in: Ann N Y Acad Sci. 2022 Apr;1510(1):180-181. PMID: 32659041.

65. Pin TW, Butler PB. The effect of interactive computer play on balance and functional abilities in children with moderate cerebral palsy: A pilot randomized study. Clin Rehabil. 2019 Apr;33(4):704-710. doi: 10.1177/0269215518821714. Epub 2019 Jan 2. PMID: 30599772.

66. Thomas SS, Buckon CE, Phillips DS, Aiona MD, Sussman MD. Interobserver reliability of the gross motor performance measure: Preliminary results. Dev Med Child Neurol. 2001 Feb;43(2):97-102. doi: 10.1017/s0012162201001000160. PMID: 11221911.

67. Jacobs JM. Management options for the child with spastic cerebral palsy. Orthop Nurs. 2001 May-Jun;20(3):53-59; quiz 59-61. doi: 10.1097/00006416-200105000-00009. PMID: 12025635.

68. Hodapp M, Vry J, Mall V, Faist M. Changes in soleus H-reflex modulation after treadmill training in children with cerebral palsy. Brain. 2009 Jan;132(Pt 1):37-44. doi: 10.1093/brain/awn287. Epub 2008 Nov 3. PMID: 18984603.

69. Wu YN, Hwang M, Ren Y, Gaebler-Spira D, Zhang LQ. Combined passive stretching and active movement rehabilitation of lower-limb impairments in children with cerebral palsy using a portable robot. Neurorehabil Neural Repair. 2011 May;25(4):378-385. doi: 10.1177/1545968310388666. Epub 2011 Feb 22. PMID: 21343525.

70. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. Phys Ther. 2003 Aug;83(8):713-721. PMID: 12882612.

71. Laver KE, Lange B, George S, Deutsch JE, Saposnik G, Crotty M. Virtual reality for stroke rehabilitation. Cochrane Database Syst Rev. 2017 Nov 20;11(11):CD008349. doi: 10.1002/14651858.CD008349.pub4. PMID: 29156493; PMCID: PMC6485957.

72. Ghai S, Driller M, Ghai I. Effects of joint stabilizers on proprioception and stability: A systematic review and meta-analysis. Phys Ther Sport. 2017 May;25:65-75. doi: 10.1016/j.ptsp.2016.05.006. Epub 2016 May 18. PMID: 28262354.

73. Ghai S, Driller MW, Masters RSW. The influence of below-knee compression garments on knee-joint proprioception. Gait Posture. 2018 Feb;60:258-261. doi: 10.1016/j.gaitpost.2016.08.008. Epub 2016 Aug 9. PMID: 27523397.

74. Ronsse R, Puttemans V, Coxon JP, Goble DJ, Wagemans J, Wenderoth N, Swinnen SP. Motor learning with augmented feedback: Modality-dependent behavioral and neural consequences. Cereb Cortex. 2011 Jun;21(6):1283-1294. doi: 10.1093/cercor/bhq209. Epub 2010 Oct 28. PMID: 21030486.

75. Biffi E, Beretta E, Cesareo A, Maghini C, Turconi AC, Reni G, Strazzer S. An immersive virtual reality platform to enhance walking ability of children with acquired brain injuries. Methods Inf Med. 2017 Mar 23;56(2):119-126. doi: 10.3414/ME16-02-0020. Epub 2017 Jan 24. PMID: 28116417.

76. Malaiya R, McNee AE, Fry NR, Eve LC, Gough M, Shortland AP. The morphology of the medial gastrocnemius in typically developing children and children with spastic hemiplegic cerebral palsy. J Electromyogr Kinesiol. 2007 Dec;17(6):657-663. doi: 10.1016/j.jelekin.2007.02.009. Epub 2007 Apr 24. PMID: 17459729.

77. Lampe R, Grassl S, Mitternacht J, Gerdesmeyer L, Gradinger R. MRT-measurements of muscle volumes of the lower extremities of youths with spastic hemiplegia caused by cerebral palsy. Brain Dev. 2006 Sep;28(8):500-506. doi: 10.1016/j.braindev.2006.02.009. Epub 2006 May 11. PMID: 16690238.

78. Wiley ME, Damiano DL. Lower-extremity strength profiles in spastic cerebral palsy. Dev Med Child Neurol. 1998 Feb;40(2):100-107. doi: 10.1111/j.1469-8749.1998.tb15369.x. PMID: 9489498.

79. Damiano DL, Abel MF. Functional outcomes of strength training in spastic cerebral palsy. Arch Phys Med Rehabil. 1998 Feb;79(2):119-125. doi: 10.1016/s0003-9993(98)90287-8. PMID: 9473991.

80. Desloovere K, Molenaers G, Feys H, Huenaerts C, Callewaert B, Van de Walle P. Do dynamic and static clinical measurements correlate with gait analysis parameters in children with cerebral palsy? Gait Posture. 2006 Nov;24(3):302-313. doi: 10.1016/j.gaitpost.2005.10.008. Epub 2005 Nov 21. PMID: 16303305.

81. Ross SA, Engsberg JR. Relationships between spasticity, strength, gait, and the GMFM-66 in persons with spastic diplegia cerebral palsy. Arch Phys Med Rehabil. 2007 Sep;88(9):1114-1120. doi: 10.1016/j.apmr.2007.06.011. PMID: 17826455.

82. Verschuren O, Ketelaar M, Takken T, Van Brussel M, Helders PJ, Gorter JW. Reliability of hand-held dynamometry and functional strength tests for the lower extremity in children with cerebral palsy. Disabil Rehabil. 2008;30(18):1358-1366. doi: 10.1080/09638280701639873. PMID: 18850351.

83. Lieber RL, Fridén J. Muscle contracture and passive mechanics in cerebral palsy. J Appl Physiol (1985). 2019 May 1;126(5):1492-1501. doi: 10.1152/japplphysiol.00278.2018. Epub 2018 Dec 20. PMID: 30571285; PMCID: PMC6589815.

84. Hoffer MM, Knoebel RT, Roberts R. Contractures in cerebral palsy. Clin Orthop Relat Res. 1987 Jun;(219):70-77. PMID: 3581586.

85. Gage JR, Novacheck TF. An update on the treatment of gait problems in cerebral palsy. J Pediatr Orthop B. 2001 Oct;10(4):265-274. PMID: 11727367.

86. Nordark. Transformational technologies in single-event neurological conditions: Applying lessons learned in stroke to cerebral palsy (August 14-15, 2008). Neurorehabil Neural Repair. 2009 Sep;23(7):747-765. doi: 10.1177/1545968309338028. PMID: 19710288.

87. Burdea GC, Cioi D, Kale A, Janes WE, Ross SA, Engsberg JR. Robotics and gaming to improve ankle strength, motor control, and function in children with cerebral palsy--a case study series. IEEE Trans Neural Syst Rehabil Eng. 2013 Mar;21(2):165-173. doi: 10.1109/TNSRE.2012.2206055. Epub 2012 Jul 3. PMID: 22773059; PMCID: PMC4228314.

88. Allen M. The SAGE encyclopedia of communication research methods. Sage Publications. 2017.

89. Fowler EG, Staudt LA, Greenberg MB, Oppenheim WL. Selective control assessment of the lower extremity (SCALE): development, validation, and interrater reliability of a clinical tool for patients with cerebral palsy. Dev Med Child Neurol. 2009 Aug;51(8):607-614. doi: 10.1111/j.1469-8749.2008.03186.x. Epub 2009 Feb 12. PMID: 19220390.

90. Jung S, Song S, Lee D, Lee K, Lee G. Effects of kinect video game training on lower extremity motor function, balance, and gait in adolescents with spastic diplegia cerebral palsy: A pilot randomized controlled trial. Dev Neurorehabil. 2021 Apr;24(3):159-165. doi: 10.1080/17518423.2020.1819458. Epub 2020 Sep 27. PMID: 32981401.

91. Curtis DJ, Bencke J, Mygind B. The effect of training in an interactive dynamic stander on ankle dorsiflexion and gross motor function in children with cerebral palsy. Dev Neurorehabil. 2014 Dec;17(6):393-397. doi: 10.3109/17518423.2013.844738. Epub 2014 May 27. PMID: 24865964.

92. Mutlu A, Livanelioglu A, Gunel MK. Reliability of goniometric measurements in children with spastic cerebral palsy. Med Sci Monit. 2007 Jul;13(7):CR323-329. PMID: 17599027.

93. Berg M, Jahnsen R, Froslie KF, Hussain A. Reliability of the pediatric evaluation of disability inventory (PEDI). Phys Occup Ther Pediatr. 2004;24(3):61-77. doi: 10.1300/j006v24n03_05. PMID: 15257969.

94. Bohannon RW, Smith MB. Interrater reliability of a modified Ashworth scale of muscle spasticity. Phys Ther. 1987 Feb;67(2):206-207. doi: 10.1093/ptj/67.2.206. PMID: 3809245.

95. Cohen J. Statistical power analysis for the behavioural sciences. Academic press. 2013.

96. Tur BS, Küçükdeveci AA, Kutlay S, Yavuzer G, Elhan AH, Tennant A. Psychometric properties of the WeeFIM in children with cerebral palsy in Turkey. Dev Med Child Neurol. 2009 Sep;51(9):732-738. doi: 10.1111/j.1469-8749.2008.03255.x. Epub 2009 Jan 29. PMID: 19207295.

97. Pae CU. Why systematic review rather than narrative review? Psychiatry Investig. 2015 Jul;12(3):417-419. doi: 10.4306/pi.2015.12.3.417. Epub 2015 Jul 6. PMID: 26207140; PMCID: PMC4504929.

98. Drucker AM, Fleming P, Chan AW. Research Techniques Made Simple: Assessing Risk of Bias in Systematic Reviews. J Invest Dermatol. 2016 Nov;136(11):e109-e114. doi: 10.1016/j.jid.2016.08.021. PMID: 27772550.

99. Georgiev DD, Georgieva I, Gong Z, Nanjappan V, Georgiev GV. Virtual reality for neurorehabilitation and cognitive enhancement. Brain Sci. 2021 Feb 11;11(2):221. doi: 10.3390/brainsci11020221. PMID: 33670277; PMCID: PMC7918687.

How to cite this article: Washington M, Owens RA, Moody JA. The Effects of Virtual Gaming on Lower Extremity Function in Children with Cerebral Palsy: A Narrative Review. J Biomed Res Environ Sci. 2022 July 18; 3(7): 771-778. doi: 10.37871/jbres1512, Article ID: JBRES1512, Available at: https://www.jelsciences.com/articles/jbres1512.pdf